

QCD Prediction of A_{TT} for Small Q_T Dimuon Production in pp and $p\bar{p}$ Collisions

Hiroyuki Kawamura*, Jiro Kodaira^{1,†} and Kazuhiro Tanaka**

*Radiation Laboratory, RIKEN, Wako 351-0198, Japan

†Theory Division, KEK, Tsukuba 305-0801, Japan

**Department of Physics, Juntendo University, Inba, Chiba 270-1695, Japan

Abstract. We present QCD prediction of double-spin asymmetries (A_{TT}) in transversely polarized Drell-Yan process at small transverse momentum Q_T of dimuon. Resummation of large logarithmic corrections, relevant in small Q_T region, is performed up to next-to-leading logarithmic (NLL) accuracy. A_{TT} at RHIC, J-PARC and GSI are studied numerically in the corresponding kinematic regions. We show that the large A_{TT} is obtained for small Q_T and moderate energies.

Keywords: Transversity, Drell-Yan process, Soft gluon resummation, Double-spin asymmetry

PACS: 12.38.-t, 12.38.Cy, 13.85.Qk, 13.88.+e

The transversity $\delta q(x)$ is one of the twist-2 distribution functions of nucleon, which represents the distribution of transversely polarized quarks inside transversely polarized nucleon [1]. Although it is theoretically as important as other twist-2 distribution functions such as the unpolarized parton distributions ($q(x), g(x)$) and the helicity distributions ($\Delta q(x), \Delta g(x)$), very little has been known of it so far. This is because $\delta q(x)$ is a chiral-odd function, and should be always accompanied with another chiral-odd function in physical observable, and therefore cannot be measured in inclusive DIS. Transversely polarized Drell-Yan (tDY) process, $p^\uparrow p^\uparrow \rightarrow \ell^+ \ell^- X$, $p^\uparrow \bar{p}^\uparrow \rightarrow \ell^+ \ell^- X$, is one of the processes where we can access the transversity by measuring the double-spin asymmetry: $A_{TT} \equiv \frac{\Delta_T d\sigma}{d\sigma} \equiv \frac{d\sigma^{\uparrow\uparrow} - d\sigma^{\uparrow\downarrow}}{d\sigma^{\uparrow\uparrow} + d\sigma^{\uparrow\downarrow}}$. Since tDY is inclusive in the final state, it is in principle the cleanest process to access $\delta q(x)$. However, at the RHIC-Spin experiments, the asymmetries are likely to be quite small [2]. This comes from the fact that the DY process at pp collider probes the sea distributions which are likely to be small for the transversity. Moreover, the rapid growth of unpolarized sea distributions enhances the denominator of A_{TT} at low- x which is typically probed by tDY at RHIC. On the other hand, much larger asymmetries are expected at the proposed spin experiments at J-PARC and GSI, which are to be performed at lower energies [3, 4, 5].

In this work, we study the asymmetries in Q_T distribution of dimuon, especially at small Q_T where the bulk of dimuon is produced. At small Q_T , the cross section is not described correctly by the fixed order calculations since large logarithmic corrections such as $\alpha_s^n \log^m(Q^2/Q_T^2)/Q_T^2$ ($m \leq 2n - 1$) appear at each order of perturbation series. These so-called “recoil logs” come from soft gluon emissions, and have to be resummed to all orders in α_s to make a reliable prediction of the cross section at small Q_T .

¹ Deceased on Sep.16, 2006.

The resummation is carried out in the impact parameter b space, conjugate to Q_T space, to enforce transverse-momentum conservation, and the resummed cross section is expressed as the Fourier transform back to the Q_T space. Here we perform the resummation at the next-to leading logarithmic (NLL) accuracy, which corresponds to adding up the terms with $m = 2n - 1$ and $2n - 2$, respectively, for all n . At this level, the “resummed part” of the spin-dependent cross section of tDY, differential in invariant mass Q , transverse momentum Q_T and rapidity y of dimuon, and in the azimuthal angle ϕ of one of the outgoing leptons is given by [7] ($\sqrt{S}$ is the CM energy of hadron system),

$$\begin{aligned} \frac{\Delta_T d\sigma^{\text{NLL}}}{dQ^2 dQ_T^2 dy d\phi} &= \cos(2\phi) \frac{\alpha^2}{3N_c S Q^2} \sum_i e_i^2 \int_0^\infty db \frac{b}{2} J_0(bQ_T) e^{S(b,Q)} \\ &\times \left[(C_{qq} \otimes \delta q_i) \left(x_1^0, \frac{b_0^2}{b^2} \right) (C_{\bar{q}\bar{q}} \otimes \delta \bar{q}_i) \left(x_2^0, \frac{b_0^2}{b^2} \right) + (x_1^0 \leftrightarrow x_2^0) \right]. \end{aligned} \quad (1)$$

Here $J_0(bQ_T)$ is a Bessel function, $b_0 = 2e^{-\gamma_E}$ with γ_E the Euler constant. The large logarithmic corrections are resummed into the Sudakov factor $e^{S(b,Q)}$ with $S(b,Q) = -\int_{b_0^2/b^2}^{Q^2} (d\kappa^2/\kappa^2) \{ \ln \frac{Q^2}{\kappa^2} A_q(\alpha_s(\kappa)) + B_q(\alpha_s(\kappa)) \}$. Perturbatively calculable quantities, i.e., A_q , B_q and the coefficient functions $C_{qq}(z)$ and $C_{\bar{q}\bar{q}}(z)$, are expressed as power series in α_s , and their explicit forms necessary at the NLL accuracy are found in Ref.[7]. $x_{1,2}^0 = (Q/\sqrt{S})e^{\pm y}$, and $\delta q_i(x, \mu^2)$ is the transversity of i -th flavour quark at the $\overline{\text{MS}}$ scale μ . The singularity in b -integration, due to Landau pole in $\alpha_s(\kappa)$, is taken care of by “contour deformation method” introduced in [8]. Correspondingly, the nonperturbative effects are included by the replacement $e^{S(b,Q)} \rightarrow e^{S(b,Q) - g_{NP} b^2}$ in (1) [6, 8, 9], with a non-perturbative parameter g_{NP} , which (roughly speaking) parameterizes the intrinsic k_T -distribution of quarks inside nucleon. Then we combine the resummed part (1), which embodies the logarithmically enhanced contributions for small Q_T to all orders, with the “residual part” of the fixed-order cross section, which is not associated with such logarithmic enhancement. To perform this consistently, we need the leading order (LO) tDY cross section at finite Q_T ; this is of $O(\alpha_s)$ and is obtained as QCD prediction at large Q_T [7]. The matching of the NLL formula (1) with the corresponding component in the LO cross section is performed at intermediate Q_T following the formulation of Ref.[9], to ensure no double counting for all Q_T , and we obtain the “NLL+LO” prediction of the tDY cross section, $\Delta_T d\sigma^{\text{NLL+LO}}/(dQ^2 dQ_T^2 dy d\phi)$, which has a uniform accuracy over the entire range of Q_T [7]. The “NLL+LO” cross section of unpolarized Drell-Yan process, $d\sigma^{\text{NLL+LO}}/(dQ^2 dQ_T^2 dy d\phi)$, is obtained in the same way, utilizing the results in the literature [6, 10].

Taking the ratio of these cross sections, we obtain the double-spin asymmetries:

$$A_{TT} = [\Delta_T d\sigma^{\text{NLL+LO}}/dQ^2 dQ_T^2 dy d\phi] / [d\sigma^{\text{NLL+LO}}/dQ^2 dQ_T^2 dy d\phi]. \quad (2)$$

In order to calculate A_{TT} numerically, we need to assume a model of the transversity $\delta q_i(x)$ for the numerator. Here we take a model used in [2], which saturates the Soffer bound [11] as $\delta q_i(x, \mu_0^2) = [q_i(x, \mu_0^2) + \Delta q_i(x, \mu_0^2)]/2$ at the low input scale $\mu_0 \sim 0.6\text{GeV}$ and is evolved to the higher μ^2 with NLO DGLAP kernel [12]; as the inputs, we




FIGURE 1. A_{TT} at NLL+LO accuracy in pp collision at RHIC kinematics: $\sqrt{S} = 200$ GeV, $Q = 3, 5, 8, 15$ GeV and $\phi = 0$ with $y = 0$ (left panel) and $y = 2$ (right panel).




FIGURE 2. A_{TT} at NLL+LO accuracy. Left: $\sqrt{S} = 10$ GeV, $Q = 2, 3, 4$ GeV, $\phi = 0$, and $y = 0$ for pp collision at J-PARC kinematics. Right: $\sqrt{S} = 14.5$ GeV, $Q = 2, 3, 4, 6$ GeV, $\phi = 0$, and $y = 0$ for $p\bar{p}$ collision at GSI kinematics.

use GRV98 [13] for $q_i(x, \mu_0^2)$ and GRSV2000 [14] for $\Delta q_i(x, \mu_0^2)$. Correspondingly, the GRV98 distributions are used for calculating the denominator of (2). The non-perturbative parameter g_{NP} are taken to be common in the numerator and the denominator of (2), and we use $g_{NP} = 0.5 \text{ GeV}^2$ which is consistent with the result of [15].²

Fig. 1 shows A_{TT} in pp collision at RHIC kinematics: $\sqrt{S} = 200$ GeV, $Q = 3, 5, 8, 15$ GeV and $\phi = 0$, with $y = 0$ (left panel) and $y = 2$ (right panel). In this case, A_{TT} are about 4-8% and rather flat in the small Q_T region where the resummed cross sections are dominant in both the numerator and denominator of (2). We have smaller A_{TT} for smaller Q due to the growth of the sea distributions in the denominator of (2) at small x . We obtain slightly larger A_{TT} for $y = 2$ compared with the $y = 0$ case, and it appears that generically the y -dependence of A_{TT} is small.

The left panel of Fig. 2 is same as Fig. 1, but at J-PARC kinematics: $\sqrt{S} = 10$ GeV,

² The g_{NP} -dependence of the polarized and unpolarized cross sections almost cancels in A_{TT} of (2) in the range $g_{NP} = 0.3\text{-}0.8 \text{ GeV}^2$.

$Q = 2, 3, 4$ GeV, $\phi = 0$ and $y = 0$. In this kinematics, the parton distributions at medium x ($x_{1,2}^0 = 0.2-0.4$) are probed, so that the transversity distributions in the numerator of (2) are larger than the RHIC case while the growth of the sea distributions in the denominator is not significant. Therefore we obtain much larger $A_{TT} \sim 15\%$ than the RHIC case of Fig. 1. A_{TT} are again flat as functions of Q_T , and the dependence on Q is also weak.

The right panel of Fig. 2 shows A_{TT} in $p\bar{p}$ collision at GSI kinematics: $\sqrt{S} = 14.5$ GeV, $Q = 2, 3, 4, 6$ GeV, $\phi = 0$ and $y = 0$, where A_{TT} are dominated by valence distributions at medium x . The largest A_{TT} of 15-30% are obtained in this case. The results are extremely flat as functions of Q_T . Integrating the numerator and the denominator of (2) over Q_T^2 , we reproduced the NLO asymmetries given by Barone et al.[3]. We also calculated the y -dependence of the results in Fig. 2, and it turns out to be small for both J-PARC and GSI cases.

To summarize, we have calculated the double-spin asymmetries A_{TT} for small Q_T DY pair production in high-energy pp collisions at RHIC and in moderate-energy pp and $p\bar{p}$ collisions at J-PARC and GSI. Our results demonstrate that A_{TT} reach a finite value at $Q_T = 0$ through the flat behavior in the small Q_T region, reflecting that the soft gluon resummation to the NLL level has universal structure for the polarized and unpolarized DY [7], and also reveal that A_{TT} are large enough to be experimentally measured, especially in moderate energies.

ACKNOWLEDGMENTS

We would like to thank Werner Vogelsang, Hiroshi Yokoya and Stefano Catani for valuable discussions and comments. The work of J.K. and K.T. was supported by the Grant-in-Aid for Scientific Research Nos. C-16540255 and C-16540266.

REFERENCES

1. J. P. Ralston and D. Soper, *Nucl. Phys.* **B152**, 109 (1979).
2. O. Martin, A. Schäfer, M. Stratmann and W. Vogelsang, *Phys. Rev.* **D57**, 3084 (1998); *ibid.* **D60**, 117502 (1999).
3. V. Barone, A. Cafarella, C. Coriano, M. Guzzi and P. G. Ratcliffe, *Phys. Lett* **B639** 483 (2006).
4. H. Shimizu, G. Sterman, W. Vogelsang and H. Yokoya, *Phys. Rev.* **D71** 114007 (2005); H. Yokoya, in these proceedings.
5. A. Drago, in these proceedings.
6. J. C. Collins, D. Soper and G. Sterman, *Nucl. Phys.* **B250** 199 (1985).
7. H. Kawamura, J. Kodaira, H. Shimizu and K. Tanaka, *Prog. Theor. Phys.* **115**, 667 (2006).
8. E. Laenen, G. Sterman and W. Vogelsang, *Phys. Rev.* **D63** 114018 (2001); S. Kulesza, G. Sterman and W. Vogelsang, *Phys. Rev.* **D66** 014001 (2002).
9. G. Bozzi, S. Catani, D. de Florian and M. Grazzini, *Phys. Lett.* **B564** 65 (2003); *Nucl. Phys.* **B737** 73 (2006).
10. G. Altarelli, R. K. Ellis, M. Greco and G. Martinelli, *Nucl. Phys.* **B246**, 12 (1984).
11. J. Soffer, *Phys. Rev. Lett.* **74** 1292 (1995).
12. A. Hayashigaki, Y. Kanazawa and Y. Koike, *Phys. Rev.* **D56** 7350 (1997); S. Kumano and M. Miyama, *Phys. Rev.* **D56** 2504 (1997); W. Vogelsang, *Phys. Rev.* **D57** 1886 (1998).
13. M. Glück, E. Reya and A. Vogt, *Eur. Phys. J.* **C5** 461 (1998).
14. M. Glück, E. Reya, S. Stratmann, and W. Vogelsang, *Phys. Rev.* **D63** 094005 (2001).
15. A. Kulesza and W. J. Stirling, *JHEP* **0312** 056 (2003).